arXiv:astro-ph/0402016v1 1 Feb 2004

Disappearance of N_2H^+ from the Gas Phase in the Class 0 Protostar IRAM 04191

A. Belloche[1,2] and P. André[3]

[1] Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany, `belloche@mpifr-bonn.mpg.de`
[2] LRA/LERMA, Ecole Normale Supérieure, 24 rue Lhomond, F-75231 Paris Cedex 05, France
[3] Service d'Astrophysique, CEA/Saclay, F-91191 Gif-sur-Yvette, France, `pandre@cea.fr`

Abstract. We present a high-resolution spectroscopic study of the envelope of the young Class 0 protostar IRAM 04191+1522 in Taurus. N_2H^+(1-0) observations with the Plateau de Bure Interferometer and the IRAM 30m telescope demonstrate that the molecular ion N_2H^+ disappears from the gas phase in the inner envelope ($r < 1600$ AU, $n_{\mathrm{H}_2} > 4-7\times10^5$ cm^{-3}). This may result from N_2 depletion on polar ice mantles and enhanced grain chemistry.

1 The Class 0 Protostar IRAM 04191

With an age $t \sim 1-3\times10^4$ yr measured from the beginning of the accretion phase, IRAM 04191+1522 is the youngest Class 0 protostar known so far in the Taurus molecular cloud (d = 140 pc). It has a prominent envelope (1.5 $M_\odot$) and an associated powerful bipolar outflow [3]. Belloche et al. [4] showed that the envelope is undergoing both extended infall motions and fast, differential rotation. They proposed that the rapidly rotating inner part of the envelope (r < 3500 AU) corresponds to a magnetically supercritical core decoupling from an environment still supported by magnetic fields and strongly affected by magnetic braking.

We recently carried out new observations in the N_2H^+(1-0) line with the Plateau de Bure Interferometer (PdBI) to probe the inner structure of the IRAM 04191 envelope. Here, we present and discuss the results of these high-resolution ($\sim 5''$) observations of the small-scale distribution of N_2H^+.

2 Small-Scale N_2H^+(1-0) Emission

Fig. 1a shows the N_2H^+(1-0) integrated intensity map obtained with PdBI, which filters out the extended emission observed with the IRAM 30m telescope. The (naturally-weighted) synthesized half-power beamwidth is $5.8'' \times 4''$. Two emission peaks are seen approximately along the direction perpendicular to the outflow axis, on either side of a gap very close to the



Fig. 1. **(a)** N_2H^+(1-0) integrated intensity map of the IRAM 04191 envelope obtained with PdBI. The contour step is 0.1 Jy beam^{-1} km s^{-1} (i.e. 3 times the rms noise). The big circle represents the primary beam. The emission gap very close to the central position (*black cross*) suggests a strong decrease of the N_2H^+ abundance in the inner part of the envelope. **(b)** Map of the LSR velocity (*squares*) as derived from Gaussian hfs fits to the N_2H^+(1-0) spectra. The map of integrated intensity is overlaid as contours. We measure a mean velocity gradient of $\sim$ 26 km s^{-1} pc^{-1} across the envelope (*black arrow*), whose direction at PA $\sim 107°$ is consistent with the rotation seen on larger scale.

central position. This suggests a strong decrease of the N_2H^+ abundance toward the center. Besides, the two emission peaks have very different centroid velocities (see Fig. 1b): we measure a strong velocity gradient of $\sim$ 26 km s^{-1} pc^{-1} along the axis with PA = 107°, i.e. approximately the same axis along which a rotational velocity gradient is detected on larger scale [4]. The gradient measured with PdBI is 8 times larger than the mean velocity gradient observed with the 30m telescope toward the core, which confirms the presence of fast, differential rotation in the IRAM 04191 envelope.

3 N_2H^+ Disappearance from the Gas Phase

Since the interferometer filters out the extended emission, we added short spacings obtained with the 30m telescope [4] to the PdBI data. As shown in the N_2H^+(1-0) integrated intensity map (see Fig. 2a), the emission is resolved in a ringlike structure with two local peaks, surrounding a gap close to the center. Gaussian hfs fits to the 7-component N_2H^+(1-0) spectra yield a mean total opacity $\tau_{\rm tot} = 10 \pm 4$ in the central part ($\theta < 10''$), corresponding to $\tau = 2.6 \pm 1.0$ for the strongest component and $\tau = 1.1 \pm 0.4$ for the isolated component. However, the opacity map does not show any enhancement toward the center and the gap close to the center does not have the



Fig. 2. **(a)** N_2H^+(1-0) integrated intensity map of the IRAM 04191 envelope combining the PdBI data and the short spacings obtained at the 30m telescope. The contour step is 0.16 Jy beam^{-1} km s^{-1} (i.e. 7 times the rms noise). The white circle represents the PdBI primary beam. **(b)** Radial profile of the N_2H^+(1-0) integrated intensity corrected for the primary beam attenuation. The dots with error bars represent the average intensity in concentric shells limited to the positions shown in the insert. The solid curve indicates the H_2 column density profile [10, 4] scaled to the 9th shell for comparison. The dashed curves show the effect of a hole of N_2H^+(1-0) emission with radius r = 1350 and 1850 AU, respectively. They suggest a strong decrease of the N_2H^+ abundance in the inner part of the envelope ($r < 1600$ AU).

highest opacity. Besides, the intensity maps integrated over each ΔF_1 group of N_2H^+(1–0) hyperfine components separately show the same pattern. We are thus confident that the observed emission gap does not result from an opacity effect.

Assuming a constant excitation temperature for N_2H^+(1-0), we compare in Fig. 2b the circularly-averaged profile of the N_2H^+(1-0) integrated intensity, corrected for PdBI primary beam attenuation, with the H_2 column density profile derived from the dust emission [10, 4]. The N_2H^+(1-0) profile departs from the H_2 column density profile for $\theta \leq 13''$. A good model fit is obtained when one assumes a hole of N_2H^+(1-0) emission in the central part of the envelope ($r <$ 1350-1850 AU, see dashed curves). This crude comparison thus strongly suggests that the N_2H^+ ion disappears from the gas phase above a density of $n_{\mathrm{H_2}} \sim 4 - 7 \times 10^5$ cm^{-3} in the IRAM 04191 envelope.

4 Discussion

Very little evidence for N_2H^+ abundance drop has been reported so far in dense cores (see however the case of B68 [6]). The N_2H^+(1-0) emission usually follows the dust emission very well in prestellar cores and has long been thought to be one of the best molecular tracers of dense gas [12, 7]. From a theoretical point of view, all recent chemical models predict a centrally-peaked N_2H^+ column density profile during the prestellar collapse phase

[5, 1, 2, 9, 11]. In particular, the models calculated by Aikawa et al. [2] including gas-phase reactions, gas-dust interactions and diffusive grain-surface reactions predict an increase of the N_2H^+ abundance during the core collapse up to $n_{H_2} \sim 1 \times 10^7$ cm^{-3} or more. Since only a few 10^4 yr have elapsed and the density at $r \sim 1600$ AU cannot have changed much between point mass formation (the last stage of this model) and the present stage of IRAM 04191, the drop observed in the N_2H^+ abundance above $n_{H_2} \sim 4 - 7 \times 10^5$ cm^{-3} is strongly at variance with current theoretical predictions.

Our result implies either a stronger dependence of the N_2H^+ abundance on density (i.e. an earlier drop in abundance during the prestellar phase) or an abundance drop on a timescale as short as a few 10^4 yr. Since N_2 is the precursor of N_2H^+, the drop in N_2H^+ abundance may result from stronger N_2 depletion onto grain surfaces. This could happen if N_2 were depleted on H_2O ice mantles characterized by a higher binding energy (see [6], Model C of [1] and Fig. 6 of [11]) and/or if the chemistry transforming N_2 in less volatile species on grain surfaces were enhanced. It is also possible that N_2H^+ destroyers such as CO have returned to the gas phase in the inner envelope, following heating by the central protostar or shock by the outflow. However, the dust temperature is below 10 K at $r \sim 1000$ AU and our $C^{18}O$(2-1) observations [4] at a resolution of 800 AU in radius show depletion toward the envelope center by a factor of 3 in column density when compared to the 1.3mm continuum emission. This latter explanation seems thus less likely.

A better tracer than N_2H^+ will thus have to be found to probe the velocity structure of cold protostellar envelopes at high resolution with the next generation of (sub)millimeter instruments such as ALMA. H_2D^+ may be such a good tracer [6, 8].

Acknowledgements. We would like to thank Y. Aikawa, E. Bergin, H. Fraser, M. Gerin, D. Johnstone, G. Pineau des Forêts and M. Walmsley for enlightening discussions on chemistry, and F. Gueth, R. Neri and J. Pety for their help with the reduction of our PdBI observations.

References

1. Y. Aikawa, N. Ohashi, S.I. Inutsuka, et al.: ApJ, **552**, 639 (2001)
2. Y. Aikawa, N. Ohashi, & E. Herbst: ApJ, **593**, 906 (2003)
3. P. André, F. Motte, & A. Bacmann: ApJ, **513**, L57 (1999)
4. A. Belloche, P. André, D. Despois, & S. Blinder: A&A, **393**, 927 (2002)
5. E.A. Bergin, & W.D. Langer: ApJ, **486**, 316 (1997)
6. E.A. Bergin, J. Alves, T.L. Huard, & C.J. Lada: ApJ, **570**, L101 (2002)
7. P. Caselli, P.J. Benson, P.C. Myers, & M. Tafalla: ApJ, **572**, 238 (2002)
8. P. Caselli et al: A&A, **403**, L37 (2003)
9. Z.Y. Li, V.I. Shematovich, D.S. Wiebe, & B.M. Shustov: ApJ, **569**, 792 (2002)
10. F. Motte, & P. André: A&A, **365**, 440 (2001)
11. V.I. Shematovich, D.S. Wiebe, B.M. Shustov, & Z.Y. Li: ApJ, **588**, 894 (2003)
12. M. Tafalla, P.C. Myers, P. Caselli, et al.: ApJ, **569**, 815 (2002)